e ` ¶-@/¢ e

B g <š;ó=l F¢ • ù G P S6= ‹ P Ï51 =l'â -+° • B-@/¢

O˜,« 'L… L) † K }TÅ E¿ ” Ç

Yî(4` ƒFe bx•|ÁÀ $ ‘` wË!ñ•P !Ra'a 44 4 4 4T4

, +° U*T=l ? e Ç *>î B g<š;ó=l F¢ U‹(# !UŒ • B % ‘ ã æ ¸ G P S6= ‹ P U‹4 "UŒÏ51 =l'â -+° • ù å" 3ß ‘ 47 4 U"Q g.ý S ’ g gKŽ b!ç+° 4 ") Ï51 · U‹ KE5 T° S W eFy+° 9 ‘ ã æ ˆ f %&“3ð U‹bKò l 5-@/¢ %A•+° ` ¶Bp Å ö=l"Þ c X 2 9 8 Û Ò"• Ò+° (# ! Ì ‘-@/¢=l"Þ c X Y UŒ =x=l ? e Ç * ì"w U‹2 ¼ P3ðK =l ? e Ç * î œ =l ù À "K U‹2 hUŒ=l Ô Ç ’ "K U‹+ hUŒ &>þU‹ >þUÇ, S [q U‹- ! hUŒÛ@Ô (# ! % 4 ")Ï51 =l'â -+° • ù 3ÿ È 4 ")Ï513ð*c · Ÿ · U•T°+° -U‹/ UŒ U‹ ^ UŒU‹ ã % &“3ð*c

2 h ¸ + h!`

ŸB¶P U• >þ ¸ - ! hB¶ z ‘ ã %&“3ð T#' Ì ^ 5 M+Ł (# ! : 2 hK +° . 54 @Í ’ ; u U‹0 UŒU‹h >þ ¸ - ! h . s 4 @Í ’ ; u U‹0 UŒ4Ž')Ï513ð T#' Ì ^ 5 M+Ł (# ! : 2 h . 4 @Í ’ ; u U‹0 UŒU‹h >þ ¸ - ! h . s 54 @Í ’ ; u U‹0 UŒ ‘ ã %&“3ð 4 ")Ï513ð ö • œ š+° 2 h : (# !#' Ì nFû t •,$ Ÿ U‹R U‹s 0 UŒU‹h : (# !#' Ì nFû t •,$ Ÿ U‹R U‹s 0 UŒU‹þ : (# !#' Ì nFû s tBK,$ Ÿ U‹R ÃQ \5 Ã\ •` $ ‘` wË!ó (1#M'¥'þ nâ , F' • Fû s tBK,$ ŸU‹R U‹s 0 UŒ3ÿ@æ (# ! 9 F ÌFù ‘ ã æ ¸ 4 ")Ï51=l"Þ +° ï 'â - U¢ @K» Ì ‘=l Y (# !"• Ò Ê Ì U‹P3ð+°=l"Þ c X s=”(Ü & &6) Ç û z+°B· ë - ŸJZ@ù 6= ‹ P U‹G P SUž B g<š;ó=l F¢ Už =l ? Už Ž=l &FýM¥, U‹) âFù%å-@ ý@Í Þ‹ 9 & # U• 9 & # UŒ

Effect of hemocoagulase agkistrodon on coagulation status in patients with traumatic brain injury in vitro

Ma Ruf[1], Ren Jing[2], Zhang Zhubo[2], Men Jianlong[2], Deng Quanjun[2]

[1]Precision Medicine Center, Tianjin Medical University General Hospital, Tianjin 300052, China; [2]Department of Neurosurgery, Tianjin Medical University General Hospital, Tianjin 300052, China

Corresponding author: Men Jianlong. Email: thromhaemo@126.com; Deng Quanjun. Email: tjmugh2020@126.com

- Abstract . Objective To analyze the effects of hemocoagulase agkistrodon (HCA) on the coagulation status of healthy people and traumatic brain injury (TBI) patients in vitro. Methods A total of 10 TBI patients were enrolled from December 2018 to June 2019, and 24 age and sex matched healthy controls were also enrolled. Clinical and pathological data, blood samples of all

?*TUŒE" 0*T!`#â? /,Fö Ž < ÊO¸ •2' ±+°"• Ò
Û ÒU•2 W f
5 MU{L#!"• Ò Û Ò î5/
,lOú+°-@/¢U¢ š • †E ,'A/ 2 L
"=l 3 X Ì 'Fy ƒ + A OŒÍ ±NUMÌ >##, 3*T =
ù À ›E< M Ì ' M O L , #AAn# L L 9 8"•
Ò+° (#! Œ‹%&"3ð Ì '" 0*T!` ŒŒ
+ A OŒÍ ±N=l 3 X Œ•ó*T4º) 4 %'ã ·
=l ? e Ç * E =x ì"w Uì"w 3 3 1 U•=l ù À
"K U‹2 hUŒ=l Ô Ç '"K U‹+ hUŒ &>þU‹ >þŒ
¸, S [q U‹-! hUŒ=l ? e Ç * Ž
5 4 @Í '0)2
Fó*T % X C E L&/÷ œ š ¿ U•3 0 3 3 D›"E
=x4 @Í '2¼ 3û 3 H A P I R ìO¸7U•2LhK + h >þ
-! h œ š s 90R 4 • - 2 / U•ó*T -U‹0 U•0 UŒ"
.f UžT° 90R 4 • - 2 / U•ó*T -U‹8/ UŒ".f Fó
*T + R U S K A L(ã ïOA!LDL¯l9S8"• Ò Û ÒK +° 2
h + h >þ -! h+° . U0,$Eç"• ÒK 6E 3 3
+° P P ! D¯ Fó*T - A N N H I T N @ ƒFe hY' ÁÀ $ '` wË óc T '` wË!óc 60 SHc " €+° P P ! D¯ Ò+° P P ! D¯ +° P

5 (#!,$Eç"• Ò 7=l ? e Ç * 3 3+°! D¯
T 2 W#'Ì ^ 5 M•LÒ+° (#!: U•%=l
? e Ç * î œE =x! D¯ U• ã %&“3ð 2 h . 54
@Í ' ; u U‹' Ý U•0 UŒEU•h >þ ¸ -! h
. 4 @Í ' ; u U‹' Ýh 2 W f ¸ U•
0 UŒEU•Ž Ï513ð+° 2 h . 4
@Í ' ; u U‹' Ý U•0 UŒEU•h >þ ¸ -! h 5
4 @Í ' ; u U‹' Ýh 2 W f ¸ U•0
UŒ=” j.f U•@æ [‘ ã %&“3ð
B [4 " Ï513ð U•T#'Ì (#!: U•TD¯ z"• Ò8/
± +° ,$ Eç œ šU‹ : 5 M L 5 M:L
5 M L 5 M:L 5 M L 5 M:L
5 MŒEU•h +h: >þ ¸ -! h"• Ò S " c 7A 1R

=lF¢ Ž <Dó/7 U•2 h4U- ü.fP 'â - U•'J« ü
.f z 'â - Už h ¸ >þ ù L=l ƒ Ž <+°EK(³
U‹3Ð4 ;÷+© ¸=l ; « T=l ƒ Ž < ",$ ¾ ^*T UŒU•
+ h Ê Ì ¸ U‹BUŒ>þ û ; ü.f3Ð4 ;÷+© ËD˜ f3Ð
4 ;÷+©+°6) Ç û z Už ! h \3Ð4 ;÷+© ¸=l ; «
P åLŽ • ù U•ù L=l ƒ f Ò!` Ÿ U• ! û ; ü.f z
'â - ¸ &=l j =
X-@/¢3ÿ È j.f U•